Abdallay, A. See Grace et al., 551.

Affolter, F. See Doane et al., 1041.

Arthur, A.P. See Doane et al.., 1041.

Arthur, A.P., and P.G. Mason. Description of the immature stages and notes on the biology of Neotephritis finalis (Loew.) (Diptera: Tephritidae), a pest of sunflowers in Saskatchewan, Canada, 729.

Arthur, A.P., and Y.M. Powell. Descriptions of the immature stages and adult reproductive systems of Athrycia cinerea (Coq.) (Diptera: Tachinidae), a native parasitoid of Mamestra configurata (Walk.) (Lepidoptera: Noctuidae), 1117.

Atkinson, T.W. See Phillips et al., 933.

Auger, M. See Régnière et al., 267.

Baan, H.E. van de. See Croft et al., 121.

See Burts et al., 219.

Barnard, J. See Eigenbrode et al., 1125.

Barron, J.R. Status of the parasite Agrypon flaveolatum (Gravenhorst) (Hymenoptera, Ichneumonidae), introduced to control the winter moth in Nova Scotia and British Columbia, 11.

Barszcz, E.S. See Stevenson and Barszcz, 921.

Behan-Pelletier, V.M. Limnozetes (Acari: Oribatida: Limnozetidae) of northeastern North America, 453. Description of Scapuleremaeus kobauensis gen.nov., sp.nov. (Acari: Oribatida: Cymbaeremaeidae) from western Canada, 507.

Bellows, T.S., Jr. See Van Driesche et al., 291.

Benton, A.H. See Galloway et al., 609.

Berisford, C.W. See Clarke et al., 853.

Bohls, R.A. See Quinn et al., 589.

Boiteau, G., and D.T. Lowery. Comparison of a yellow form of the green peach aphid, Myzus persicae (Sulzer), and a green form of the tobacco aphid, Myzus nicotianae Blackman, coexisting on greenhouse potato in New Brunswick, 1029.

Boivin, G. See Michaud et al., 803.

See Sohati et al., 1127.

Borden, J.H. See Thistlewood et al., 1.

See Thistlewood et al., 737.

See Chen and Borden, 1059.

squet, Y. Descriptions of the larvae of Pterostichus ohionis Csiki and P. blanchardi Horn with a key to larvae of eastern North American Pterostichini (Coleoptera: Carabidae), 27.

Broadbent, A.B., R.G. Foottit, and G.D. Murphy. Sweetpotato whitefly Bemisia tabaci (Gennadius) (Homoptera: Aleyrodidae), a potential insect pest in Canada, 1027.

Broadbent, A.B., and D.J. Pree. Resistance to pyrazophos in the serpentine leafminer Liriomyza trifolii (Burgess) (Diptera: Agromyzidae) in Ontario greenhouses, 47.

Burts, E.C. See Croft et al., 121.

Burts, E.C., H.E. van de Baan, and B.A. Croft. Pyrethroid resistance in pear psylla, Psylla pyricola Foerster (Homoptera: Psyllidae), and synergism of pyrethroids with piperonyl butoxide, 219.

Butts, R.A. See Jones et al., 623.

Byers, J.R. Trapping male moths of Grammia blakei (Grt.) (Lepidoptera: Arctiidae) with female-baited traps at ground level, 105.

See Struble et al., 557. See Jones et al., 623.

Cameron, E.A., and R.W. Stark. Variations in the life cycle of the lodgepole terminal weevil, Pissodes terminalis Hopping (Coleoptera: Curculionidae), in California, 793.

Chen, N.-M., and J.H. Borden. Adverse effect of fenoxycarb on reproduction by the California fivespined ips, Ips paraconfusus Lanier (Coleoptera: Scolytidae), 1059.

Caénier, J.V.R., and B.J.R. Philogène. Evaluation of three trap designs for the capture of conifer-feeding beetles and other forest Coleoptera, 159.

Cigliano, M.M., R.A. Ronderos, and W.P. Kemp. Revision of the genus Elasmoderus Saussure (Orthoptera: Tristiridae), 225.

Clarke, S.R., G.L. DeBarr, and C.W. Berisford. The life history of Toumeyella pini (King) (Hymenoptera: Coccidae) in loblolly pine seed orchards in Georgia, 853.

Clouâtre, A., D. Coderre, and D. Gagnon. Habitat of a new Mymarommatidae found in southern Quebec. Canada (Hymenoptera: Terebrantes), 825.

Cloutier, C. See Maltais et al., 745.

Coderre, D. See Clouâtre et al., 825.

Cole, K.J. See Pree et al., 771.

Cooper, B.E. See Cumming and Cooper, 565.

Crawford, H.S., Jr. See Jennings and Crawford, 827.

Croft, B.A. See Burts et al., 219.

Abdallay, A. See Grace et al., 551.

Affolter, F. See Doane et al., 1041.

Arthur, A.P. See Doane et al.., 1041.

Arthur, A.P., and P.G. Mason. Description of the immature stages and notes on the biology of Neotephritis finalis (Loew.) (Diptera: Tephritidae), a pest of sunflowers in Saskatchewan, Canada, 729.

Arthur, A.P., and Y.M. Powell. Descriptions of the immature stages and adult reproductive systems of Athrycia cinerea (Coq.) (Diptera: Tachinidae), a native parasitoid of Mamestra configurata (Walk.) (Lepidoptera: Noctuidae), 1117.

Atkinson, T.W. See Phillips et al., 933.

Auger, M. See Régnière et al., 267.

Baan, H.E. van de. See Croft et al., 121.

See Burts et al., 219.

Barnard, J. See Eigenbrode et al., 1125.

Barron, J.R. Status of the parasite Agrypon flaveolatum (Gravenhorst) (Hymenoptera, Ichneumonidae), introduced to control the winter moth in Nova Scotia and British Columbia, 11.

Barszcz, E.S. See Stevenson and Barszcz, 921.

Behan-Pelletier, V.M. Limnozetes (Acari: Oribatida: Limnozetidae) of northeastern North America, 453. Description of Scapuleremaeus kobauensis gen.nov., sp.nov. (Acari: Oribatida: Cymbaeremaeidae) from western Canada, 507.

Bellows, T.S., Jr. See Van Driesche et al., 291.

Benton, A.H. See Galloway et al., 609.

Berisford, C.W. See Clarke et al., 853.

Bohls, R.A. See Quinn et al., 589.

Boiteau, G., and D.T. Lowery. Comparison of a yellow form of the green peach aphid, Myzus persicae (Sulzer), and a green form of the tobacco aphid, Myzus nicotianae Blackman, coexisting on greenhouse potato in New Brunswick, 1029.

Boivin, G. See Michaud et al., 803.

See Sohati et al., 1127.

Borden, J.H. See Thistlewood et al., 1.

See Thistlewood et al., 737.

See Chen and Borden, 1059.

squet, Y. Descriptions of the larvae of Pterostichus ohionis Csiki and P. blanchardi Horn with a key to larvae of eastern North American Pterostichini (Coleoptera: Carabidae), 27.

Broadbent, A.B., R.G. Foottit, and G.D. Murphy. Sweetpotato whitefly Bemisia tabaci (Gennadius) (Homoptera: Aleyrodidae), a potential insect pest in Canada, 1027.

Broadbent, A.B., and D.J. Pree. Resistance to pyrazophos in the serpentine leafminer Liriomyza trifolii (Burgess) (Diptera: Agromyzidae) in Ontario greenhouses, 47.

Burts, E.C. See Croft et al., 121.

Burts, E.C., H.E. van de Baan, and B.A. Croft. Pyrethroid resistance in pear psylla, Psylla pyricola Foerster (Homoptera: Psyllidae), and synergism of pyrethroids with piperonyl butoxide, 219.

Butts, R.A. See Jones et al., 623.

Byers, J.R. Trapping male moths of Grammia blakei (Grt.) (Lepidoptera: Arctiidae) with female-baited traps at ground level, 105.

See Struble et al., 557. See Jones et al., 623.

Cameron, E.A., and R.W. Stark. Variations in the life cycle of the lodgepole terminal weevil, Pissodes terminalis Hopping (Coleoptera: Curculionidae), in California, 793.

Chen, N.-M., and J.H. Borden. Adverse effect of fenoxycarb on reproduction by the California fivespined ips, Ips paraconfusus Lanier (Coleoptera: Scolytidae), 1059.

Caénier, J.V.R., and B.J.R. Philogène. Evaluation of three trap designs for the capture of conifer-feeding beetles and other forest Coleoptera, 159.

Cigliano, M.M., R.A. Ronderos, and W.P. Kemp. Revision of the genus Elasmoderus Saussure (Orthoptera: Tristiridae), 225.

Clarke, S.R., G.L. DeBarr, and C.W. Berisford. The life history of Toumeyella pini (King) (Hymenoptera: Coccidae) in loblolly pine seed orchards in Georgia, 853.

Clouâtre, A., D. Coderre, and D. Gagnon. Habitat of a new Mymarommatidae found in southern Quebec. Canada (Hymenoptera: Terebrantes), 825.

Cloutier, C. See Maltais et al., 745.

Coderre, D. See Clouâtre et al., 825.

Cole, K.J. See Pree et al., 771.

Cooper, B.E. See Cumming and Cooper, 565.

Crawford, H.S., Jr. See Jennings and Crawford, 827.

Croft, B.A. See Burts et al., 219.

Croft, B.A., E.C. Burts, H.E. van de Baan, P.H. Westigard, and H. Riedl. Local and regional resistance to fenvalerate in *Psylla pyricola* Foerster (Homoptera: Psyllidae) in western North America, 121.

Cumming, J.M., and B.E. Cooper. The identity of Micrempis anatolica Chillcott and M. bomboxynon Chillcott (Diptera: Empididae), with remarks on allied Nearctic species, 565.

Cunningham, J.C. See Otvos et al., 209.

Danks, H.V., and R.G. Foottit. Insects of the boreal zone of Canada, 625.

DeBarr, G.L. See Clarke et al., 853.

DeClerck, R. See Peschken et al., 781.

DeClerck-Floate, R. See Doane et al., 1041.

Dennehy, T.J. See Hoffman and Dennehy, 325.

Dennis, B. See Kemp and Dennis, 363.

Derby, J.L. See Peschken et al., 781.

Doane, J.F., R. DeClerck-Floate, A.P. Arthur, and F. Affolter. Description of the life stages of Macroglenes penetrans (Kirby) (Hymenoptera: Chalcidoidea, Pteromalidae), a parasitoid of the wheat midge, Sitodiplosis mosellana (Géhin) (Diptera: Cecidomyiidae). 1041.

Dosdall, L.M., and D.M. Lehmkuhl. Drift of aquatic insects following methoxychlor treatment of the Saskatchewan River system, 1077.

Duncan, R., and E.E. Lindquist. An unusually copious production of webbing by a willow-inhabiting spider mite, Schizotetranychus schizopus (Zacher), 1037.

Eigenbrode, S.D., J. Barnard, and A.M. Shelton. A system for quantifying behavior of neonate caterpillars and other small, slow-moving animals, 1125.

Elliott, N.C., and R.W. Kieckhefer. Effects of constant and fluctuating temperatures on immature development and age-specific life tables of Rhopalosiphum padi (L.) (Homoptera: Aphididae), 131.

Fairchild, W.L., and G.B. Wiggins. Immature stages and biology of the North American caddisfly genus Phanocelia Banks (Trichoptera: Limnephilidae), 515.

Farr, K.R. See Grace et al., 551.

Ferro, D.N. See Van Driesche et al., 291.

Fisher, P.A. See Pree et al., 771.

Fogal, W.H. See Grant et al., 691.

Foltz, J.L. See Phillips et al., 933.

Foottit, R.G. See Danks and Foottit, 625.

See Broadbent et al., 1027. Foster, R.N. See Quinn et al., 589.

Furniss, M.M., and J.B. Johnson. Description of the gallery and larva of *Dendroctonus punctatus* LeConte (Coleoptera: Scolytidae), 757.

Gagnon, D. See Clouâtre et al., 825.

Galloway, T.D. See Tomalak et al., 821.

Galloway, T.D., G.E. Haas, and A.H. Benton. Corrections and additions to G.P. Holland's "Fleas of Canada, Alaska and Greenland (Siphonaptera)", 609.

Gerber, G.H. The red turnip beetle, Entomoscelis americana (Coleoptera: Chrysomelidae): distribution, temperature adaptations, and zoogeography, 315.

Gooding, R.H., B.M. Rolseth, and S.A. Tarimo Nesbitt. Mapping four loci in Glossina morsitans submorsitans Newstead (Diptera: Glossinidae), 823.

Gordon, R. See Mulye and Gordon, 1111.

Gorley, R.A. See Stock and Gorley, 521.

Grace, J.K., A. Abdallay, and K.R. Farr. Eastern subterranean termite (Isoptera: Rhinotermitidae) foraging territories and populations in Toronto, 551.

Grant, G.G., W.H. Fogal, R.J. West, K.N. Slessor, and G.E. Miller. A sex attractant for the spruce seed moth, Cydia strobilella (L.), and the effect of lure dosage and trap height on capture of male moths, 691. Gray, T.G. See Struble et al., 557.

Groot, P. de. See Helson et al., 81.

Guppy, J.C., and F. Meloche. Note on Cyrtogaster vulgaris Walker (Hymenoptera: Pteromalidae), a secondary parasite of the alfalfa blotch leafminer (Diptera: Agromyzidae), with descriptions of the immature stages, 929.

Haack, R.A. See Mattson et al., 93.

Haas, G.E. See Galloway et al., 609.

Hagino, Y. Two new species of the family Brachypauropodidae (Pauropoda) from Japan, 301.

Hagley, E.A.C. Release of Chrysoperla carnea Stephens (Neuroptera: Chrysopidae) for control of the green apple aphid, Aphis pomi DeGeer (Homoptera: Aphididae), 309.

Halbert, S.E. See Robinson and Halbert, 525.

Hanson, P.E. See Miller and Hanson, 425.

Harper, P.P., and M. Lauzon. Life cycle of the nymph fly Palaeodipteron walkeri Ide 1965 (Diptera: Nymphomyiidae) in the White Mountains of southern Québec, 603.

Harris, J.L. See Jones et al., 623.

Harvey, M.S. See Smith and Harvey, 283.

Hazzard, R. See Van Driesche et al., 291.

Hébert, C. See Maltais et al., 745.

Helson, B.V., P. de Groot, J.J. Turgeon, and E.G. Kettela. Toxicity of insecticides to first-instar larvae of the spruce budmoth, Zeiraphera canadesis Mut. and Free. (Lepidoptera:Tortricidae): laboratory and field studies, 81.

Herms, D.A. See Mattson et al., 93.

Hoffman, C.J., and T.J. Dennehy. Phenology, movement, and within-field distribution of the grape berry moth, Endopiza viteana (Clemens) (Lepidoptera: Tortricidae), in New York vineyards, 325.

Hokkanen, H.M.T., and D. Pimentel. New associations in biological control: theory and practice, 829.

Holliday, N.J. See Senanayake and Holliday, 253.

Horton, D.R. Performance of a willow-feeding beetle, Chrysomela knabi Brown, as affected by host species and dietary moisture, 777.

Jaques, R.P., D.R. Laing, and H.E.L. Maw. Efficacy of mixtures of microbial insecticides and permethrin against the cabbage looper (Lepidoptera: Noctuidae) and the imported cabbageworm (Lepidoptera: Pieridae), 809.

Jennings, D.T., H.S. Crawford, Jr. Predation by red squirrels on the spruce budworm Choristoneura fumiferana (Clem.) (Lepidoptera: Tortricidae), 827.

Johnson, D.L. Spatial autocorrelation, spatial modeling, and improvements in grasshopper survey methodology, 579.

Johnson, J.B. See Furniss and Johnson, 757.

Jones, R.L. See Swedenborg et al., 245.

Jones, J.W., J.R. Byers, R.A. Butts, and J.L. Harris. A new pest in Canada: Russian wheat aphid, Diuraphis noxia (Mordvilko) (Homoptera: Aphididae), 623.

Kaupp, W.J. See Otvos et al., 209.

Kemp, W.P. See Cigliano et al., 225.

Kemp, W.P., and B. Dennis. Development of two rangeland grasshoppers at constant temperatures: development thresholds revisited, 363.

Kepner, R.L. See Quinn et al., 589.

Kettela, E.G. See Helson et al., 81.

Kieckhefer, R.W. See Elliott and Kieckhefer, 131.

Laing, D.R. See Jaques et al., 809.

Laing, J.E. See Wang and Laing, 65.

Lancisni, C.A., and B.P. Smith. Constancy of stylostome form in two water mite species, 439.

Landry, B. Geina sheppardi, a new species of grape-feeding Platyptillinae (Lepidoptera: Pterophoridae) from eastern North America, with notes on related species, 763.

Langor, D.W. Host effects on the phenology, development, and mortality of field populations of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae), 149.

Larson, D.J. Revision of North American Agabus Leach (Coleoptera: Dytiscidae): introduction, key to species groups, and classification of the ambiguus-, tristis-, and arcticus-groups, 861.

Lauzon, M. See Harper and Lauzon, 603.

Lawrence, R.K. See Mattson et al., 93.

Leech, R., and B.J. MacDonald-Leech. Dipalta serpentina Osten Sacken (Diptera: Bombyliidae) as a predator on pupae of the antlion Myrmeleon immaculatus DeGeer (Neuroptera: Myrmeleontidae) in California, 727.
Lehmkuhl, D.M. See Dosdall and Lehmkuhl, 1077.

Lindquie E.E. See Duncan and Lindquist, 1037.

Lockw J.A. Taxonomic status of the Rocky Mountain locust: morphometric comparisons of Melanoplus spre. as (Walsh) with solitary and migratory Melanoplus sanguinipes (F.), 1103.

Lowery, D.T. See Boiteau and Lowery, 1029.

Lysyk, T.J. See Régnière et al., 267.

A multiple-cohort model for simulating jack pine budworm (Lepidoptera: Tortricidae) development under variable temperature conditions, 373.

MacDonald-Leech, B.J. See Leech and MacDonald-Leech, 727.

Maher, A. See Van Driesche et al., 291.

Maltaís, J., J. Régnière, C. Cloutier, C. Hébert, and D.F. Perry. Seasonal biology of Meteorus trachynotus Vier. (Hymenoptera: Braconidae) and of its overwintering host Choristoneura rosaceana (Harr.) (Lepidoptera: Tortricidae), 745.

Mason, P.G. See Arthur and Mason, 729.

Mattson, W.J., R.A. Haack, R.K. Lawrence, and D.A. Herms. Do balsam twig aphids (Homoptera: Aphididae) lower tree susceptibility of spruce budworm?, 93.

Maw, H.E.L. See Jaques et al., 809.

McClay, A.S. See Peschken et al., 781.

McLean, J.A. See Salom and McLean, 109.

McMullen, R.D. See Thistlewood et al., 1.

See Thistlewood et al., 737.

Meloche, F. See Guppy and Meloche, 929.

Michaud, O., R.K. Stewart, and G. Boivin. Economic injury levels and economic thresholds for the green apple bug, Lygocoris communis (Knight) (Hemi, tera: Miridae), in Quebec apple orchards, 803.

Miller, G.E. See Sweeney and Miller, 569.

See Grant et al., 691.

Miller, J.C., and P.E. Hanson. Laboratory studies on development of gypsy moth, Lymantria dispar (L.) (Lepidoptera: Lymantriidae), larvae on foliage of gymnosperms, 425.

Mulye, H., and R. Gordon. Effects of selected juvenile hormone analogs on sixth-instar larvae of the eastern spruce budworm, Choristoneura fumiferana Clemens (Lepidoptera: Tortricidae), 1111.

Murphy, G.D. See Broadbent et al., 1027.

Naranjo, S.E., and A.J. Sawyer. A simulation model of northern corn rootworm, Diabrotica barberi Smith and Lawrence (Coleoptera: Chrysomelidae), population dynamics and oviposition: significance of host plant phenology, 169.

Analysis of a simulation model of northern corn rootworm, Diabrotica barberi Smith and Lawrence (Coleoptera: Chrysomelidae), dynamics in field corn, with implications for population management, 193.

Otvos, I.S., J.C. Cunningham, and W.J. Kaupp. Aerial application of two baculoviruses against the western spruce budworm, Choristoneura occidentalis Freeman (Lepidoptera: Tortricidae), in British Columbia, 209. Owen, R.E. See Packer and Owen, 431.

See Packer and Owen, 1049.

Packer, L., and R.E. Owen. Notes on the biology of Lasioglossum (Evylaeus) cooleyi (Crawford), an eusocial halictine bee (Hymenoptera: Halictidae), 431.

Allozyme variation in Halictus rubicundus (Christ): a primitively social halictine bee (Hymenoptera: Halictidae), 1049.

Perry, D.F. See Maltais et al., 745.

Peschken, D.P., A.S. McClay, J.L. Derby, and R. DeClerck. Cystiphora sonchi (Bremi) (Diptera: Cecidomyiidae), a new biological control agent established on the weed perennial sow-thistle (Sonchus arvensis L.) (Compositae) in Canada, 781.

Phillips, T.W., T.W. Atkinson, and J.L. Foltz. Pheromone-based aggregation in Orthotomicus caelatus (Eichhoff) (Coleoptera: Scolytidae), 933.

Philogène, B.J.R. See Chénier and Philogène, 159.

Pierce, H.D., Jr. See Thistlewood et al., 737.

Pimentel, D. See Hokkanen and Pimentel, 829.

Pooler, P.D. See Quinn et al., 589.

Powell, Y.M. See Arthur and Powell, 1117.

Pree, D.J. See Broadbent and Pree, 47.

Pree, D.J., K.J. Cole, and P.A. Fisher. Comparison of leaf disc and Petri dish assays for the assessment of dicofol resistance in populations of European red mite from southern Ontario, 771.

Preisler, H.K. See Robertson et al., 75.

Quicke, D.L.J., and M.J. Sharkey. A key to and notes on the genera of Braconinae (Hymenoptera: Braconidae)

from America north of Mexico with descriptions of two new genera and three new species, 337.

Quinn, M.A., R.L. Kepner, D.D. Walgenbach, R.A. Bohls, P.D. Pooler, R.N. Foster, K.C. Reuter, and J.L. Swain. Immediate and 2nd-year effects of insecticide spray and bait treatments on populations of rangeland grasshoppers, 589.

Rank, F.P. See Rank and Rank, 141.

Rank, G.H., and F.P. Rank. Diapause intensity in a French univoltine and a Saskatchewan commercial strain of Megachile rotundata (Fab.), 141.

Raske, A.G. See West et al., 55.

Régnière, J. See Maltais et al., 745.

Régnière, J., T.J. Lysyk, and M. Auger. Population density estimation of spruce budworm, Choristoneura fumiferana (Clem.) (Lepidoptera: Tortricidae), on balsam fir and white spruce from 45-cm mid-crown branch tips, 267.

Resh, V.H., and D.M. Rosenberg. Invitation paper (C.P. Alexander Fund): Spatial-temporal variability and the study of aquatic insects, 941.

Reuter, K.C. See Quinn et al., 589.

Riedl, H. See Croft et al., 121.

Robertson, J.L., S.P. Worner, and H.K. Preisler. Comparative optimal times of application of benzoylphenyl ureas to western spruce budworm, Choristoneura occidentalis Freeman (Lepidoptera: Tortricidae), 75.

Robinson, A.G., and S.E. Halbert. Review of Obsusicauda (Homoptera: Aphididae) of America north of Mexico, with a key to species, 525.

Rolseth, B.M. See Gooding et al., 823.

Ronderos, R.A. See Cigliano et al., 225.

Rosenberg, D.M. See Resh and Rosenberg, 941.

Ruberson, J.R., M.J. Tauber, C.A. Tauber, and W.M. Tingey. Interactions at three trophic levels: Edovum puttleri Grissell (Hymenoptera: Eulophidae), the Colorado potato beetle, and insect-resistant potatoes, 841.

Ryker, L.C. See Swedenborg et al., 245.

Salom, S.M., and J.A. McLean. Influence of wind on the spring flight of *Trypodendron lineatum* (Olivier) (Coleoptera: Scolytidae) in a second-growth coniferous forest, 109.

Senanayake, D.G., and N.J. Holliday. Seasonal abundance of foliage-dwelling insect pests in commercial fields and insecticide-free plots of potato in Manitoba, 253.

Sharkey, M.J. See Quicke and Sharkey, 337.

Shea, P.J. Phytophagous insect complex associated with cones of white fir, Abies concolor (Gord. and Glend.) Lindl., and its impact on seed production, 699.

Shelton, A.M. See Eigenbrode et al., 1125.

Shepherd, R.F. See Struble et al., 557.

Slessor, K.N. See Grant et al., 691.

Smith, B.P. See Lanciani and Smith, 439.

A description of the larva of Rhyncholimnochares kittatinniana Habeeb (Hydrachnidia: Limnocharidae), 445.

Smith, I.M. Description of deutonymphs and adults of Oregonacarus rivulicolus gen.nov., sp.nov. (Acari: Lebertioidea: Anisitsiellidae), 533.

North American water mites of the family Momoniidae Viets (Acari: Arrenuroidea). I. Description of adults of Cyclomomonia andrewi gen.nov., sp.nov., and key to world genera and subgenera, 543.

Description of two new species of *Platyhadracarus* gen.nov. from western North America, with remarks on classification of Athienemanniidae (Acari: Parasitengona: Arrenuroidea), 709.

North American water mites of the family Momoniidae Viets (Acari: Arrenuroidea). II. Revision of species of Momonia Halbert, 1906, 965.

North American water mites of the family Momoniidae Viets (Acari: Arrenuroidea). III. Revision of species of Stygomomonia Szalay, 1943, subgenus Allomomonia Cook, 1968, 989.

Smith, I.M., and M.S. Harvey. Description of adults of Arrenurus (?Micruracarus) kitchingi sp.nov. (Acarina: Arrenuridae) from water-filled tree holes in Australia, 283.

Smith, R.F. See Thistlewood et al., 737.

Sohati, P.H., R.K. Stewart, and G. Boivin. Egg parasitoids of the tarnished plant bug, Lygus lineolaris (P. de B.) (Hemiptera: Miridae) in Quebec, 1127.

Stark, R.W. See Cameron and Stark, 793.

Stevenson, A.B., and E.S. Barszcz. Diapause status of the overwintering generation of the carrot rust fly, Psila rosae (Fab.) (Diptera: Psilidae), in Ontario, 921.

Stewart, J.G., and L.S. Thompson. The spatial distribution of spring and summer populations of adult potato flea beetles, Epitrix cucumeris (Harris) (Coleoptera: Chrysomelidae) on small plots of potatoes, 1097.

Stewart, R.K. See Michaud et al., 803. See Sohati et al., 1127.

Stock, A.J., and R.A. Corley. Observations on a trial of broadcast burning to control an infestation of the mountain pine beetle *Dendroctonus ponderosae*, 521.

Struble, D.L., J.R. Byers, R.F. Shepherd, and T.G. Gray. Identification of sex pheromone components of the black army cutworm, *Actebia fennica* (Tauscher) (Lepidoptera: Noctuidae), and a sex attractant blend for adult males, 557.

Sundaram, A. See West et al., 55.

Swain, J.L. See Quinn et al., 589.

Sweeney, J.D., and G.E. Miller. Distribution of Barbara colfaxiana (Kearfott) (Lepidoptera: Tortricidae) eggs within and among Douglas-fir crowns and methods for estimating egg densities, 569.

Swedenborg, P.D., R.L. Jones, and L.C. Ryker. Stridulation and associated behavior of the native elm bark beetle Hylurgopinus rufipes (Eichhoff) (Coleoptera: Scolytidae), 245.

Tarimo Nesbitt, S.A. See Gooding et al., 823.

Tauber, C.A. See Ruberson et al., 841.

Tauber, M.J. See Ruberson et al., 841.

Teal, P.E.A., and J.H. Tumlinson. Neurohormonal induction of pheromone biosynthesis by *Heliothis zea* (Boddie) during the photophase, 43.

Thistlewood, H.M.A., J.H. Borden, R.F. Smith, H.D. Pierce, Jr., and R.D. McMullen. Evidence for a sex pheromone in the mullein bug, Campylomma verbasci (Meyer) (Heteroptera: Miridae), 737.

Thistlewood, H.M.A., R.D. McMullen, and J.H. Borden. Damage and economic injury levels of the mullein bug, Campylomma verbasci (Meyer) (Heteroptera: Miridae), on apple in the Okanagan Valley, 1.

Thompson, L.S. See Stewart and Thompson, 1097.

Tingey, W.M. See Ruberson et al., 841.

Togashi, I. A new sawfly, Pristiphora harai (Hymenoptera: Tenthiedinidae), collected from red spruce, Picea glehnii Masters, in Japan, 619.

Tomalak, M., H.E. Welch, and T.D. Galloway. Effects of crowding on Sulphuretylenchus spp. (Nematoda: Allantonematidae) in the haemocoel of their bark beetle hosts (Coleoptera: Scolytidae), 821.

Tumlinson, J.H. See Teal and Tumlinson, 43.

Turgeon, J.J. See Helson et al., 81.

Van Driesche, R.G., T.S. Bellows, Jr., D.N. Ferro, R. Hazzard, and A. Maher. Estimating stage survival from recruitment and density data, with reference to egg mortality in the Colorado potato beetle, *Leptin-otarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), 291.

Vickery, V.R. The biogeography of Canadian Grylloptera and Orthoptera, 389.

Walgenbach, D.D. See Quinn et al., 589.

Wang, T., and J.E. Laing. Diapause termination and morphogenesis of Holcothorax testaceipes Ratzeburg (Hymenoptera: Encyrtidae), an introduced parasitoid of the spotted tentiform leafminer, Phyllonorycter blancardella (F.) (Lepidoptera: Gracillariidae), 65.

Welch, H.E. See Tomalak et al., 821.

Weseloh, R.M. Simulation of predation by ants based on direct observations of attacks on gypsy moth larvae, 1069.

West, R.J. See Grant et al., 691.

West, R.J., A.G. Raske, and A. Sundaram. Efficacy of oil-based formulations of Bacillus thuringiensis Berliner var. kurstaki against the hemlock looper, Lambdina fiscellaria fiscellaria (Guen.) (Lepidoptera: Geometridae), 55.

Westigard, P.H. See Croft et al., 121.

Wiggins, G.B. See Fairchild and Wiggins, 515.

Worner, S.P. See Robertson et al., 75.